

LITERATURE SYNTHESIS

A literature search is a crucial component of research. It allows for the dissemination and attainment of credible data. Acquiring data must be done diligently and with credible sources. The University of Florida library and search website were used to gather information to answer the PICOT question. The literature search used CINAHL, EBSCOhost Academic Search Premier, and PubMed. CINAHL and PubMed each of which includes ~~are~~ internationally accepted and broadly accessible resource data (Ossom & Minter, 2019). When gathering data, researchers used MeSH terms to facilitate the process. MeSH stands for Medical Subject Headings and is used for "indexing journals, cataloging and searching for articles and books" (Dhammi & Kumar, 2014).

The keywords used to gather information for the PICOT question and facilitate and generate relevant information were opioid overdose, COWS screening tools, opioid emergency, National-national opioid epidemic, and opioid drug overdose. MeSH terms used to simplify the assemblage process include substance dependence, substance-related disorder, substance addiction, chemical dependence, organic mental disorders, and substance-induced. The information gathered combined meta-analysis, systemic reviews, and random control trials. The literature review was ciphered using various resources and studies, which allows for the entire diffusion of information and decreases the risk of prejudice and bias. The Academic Search Premier, CINAHL, and PubMed search engines from the UF Database provided over fifteen results when the MeSH terms of Clinical Opiate Withdrawal Scale (COWS) were used. The results from the database were further narrowed when selecting the inclusion criteria

Editing Example


publication date of five years, full text, and meta-analysis, and the results decreased to a few for each search engine. A total of six articles were selected, and they included randomized clinical trials, literature reviews, and cross-sectional studies. These studies emphasized the massive problem created in the United States and other Western countries ~~have~~ because of the opioid epidemic. Opioid disorders affect over 15 million individuals around the world. In the United States, there are over two million opioid-addicted citizens. The opioid epidemic kills over 120 thousand people annually (Dydyk et al., 2022).

The studies were selected based on their relevancy and level of evidence based on the hierarchy. There are seven accepted levels in evaluating literature in the United States. The accepted seven levels of evidence are assigned to studies based on the methodology, validity, and application to patient care. Level I is evidenced from systematic reviews and meta-analysis, level II is obtained from at least one random controlled trial, level III is obtained from a well-controlled trial without randomization, and level IV is obtained from a well-designed case or cohort study. Level V grade is obtained from systematic review qualitative studies; ~~level VI is evidence from a single, descriptive, or qualitative study; and~~ level VII is obtained from evidence from opinions of authorities or experts. For the literature review, the objective was to acquire one or two articles from different levels of evidence to compare the type of evidence of the critical nature of opioid abuse in the country (Brunt & Morris, 2023).

The opioid crisis has become a primary public health concern in recent years, with opioid withdrawal being posing a significant challenge for individuals seeking recovery. Understanding and assessing opioid withdrawal symptoms is crucial for appropriate treatment and management. This literature review aims to provide an overview of the current state of knowledge on opioid withdrawal assessment, focusing on the scales and technologies used and

Editing Example

the etiology, time course, and relationship to clinical outcomes.



Introduction

The quest to alleviate the profound challenges associated with opioid dependence and withdrawal has prompted extensive research into innovative assessment tools and therapeutic interventions. Among these endeavors, the Clinical Opiate Withdrawal Scale (COWS) has garnered significant attention for its potential to enhance patient outcomes and quality of life within skilled nursing and rehabilitation facilities. This narrative literature review delves into the existing body of research ~~surrounding~~ supporting the efficacy of COWS in improving mental health, physical stability, and social relationships for patients aged 60-98 undergoing opioid withdrawal management.

The urgency of this inquiry is underscored by the devastating impact of the opioid epidemic, which has ravaged communities across the globe. As healthcare professionals grapple with this crisis, the need for evidence-based practices and patient-centric approaches has become paramount. By exploring the fundamental mechanisms and historical context of COWS, as well as empirical evidence supporting its effectiveness, this review aims to provide a comprehensive understanding of the tool's application and implications within specialized care settings. Central to this investigation is an in-depth analysis of the literature examining COWS's influence on the key dimensions of patient well-being, including mental health, physical stability, and social engagement. ~~Studies such as Kosten & Baxter (2019) have delved into the intricate interplay between opioid withdrawal symptom management and emotional states, such as anxiety and depression. Similarly, research has illuminated the critical role of COWS in mitigating physical discomfort, a pivotal factor in facilitating rehabilitation and overall physical recovery (Dunn et al., 2020).~~

Furthermore, this review explores the literature on the impact of COWS on patients' ability to strengthen social connections and participate in meaningful relationships during their

Formatted: Right: 0.31"

Formatted: Strikethrough

Commented [JL1]: I would move this out of the intro

Formatted: Strikethrough

Formatted: Strikethrough

Editing Example

journey to recovery. As opioid withdrawal can strain interpersonal dynamics, the effective management of symptoms through tools like COWS may pave the way for improved social functioning and reintegration (Nuamah et al., 2019). ~~By synthesizing these diverse strands of inquiry, this review presents a holistic perspective on the multifaceted benefits of COWS in enhancing patient quality of life.~~

Formatted: Strikethrough

- ~~1. A comparative analysis contextualizes the findings, contrasting COWS with alternative opioid withdrawal assessment scales and illuminating their respective strengths and limitations.~~
- ~~2. Moreover, the challenges and constraints inherent in current research are critically examined, paving the way for recommendations on future avenues of investigation.~~
- ~~3. Ultimately, this comprehensive review seeks to equip healthcare professionals, policymakers, and researchers with a profound understanding of COWS's role in optimizing patient care and informing evidence-based practices within skilled nursing and rehabilitation facilities.~~

Commented [JL2]: Why are these numbered rather than integrated into the paragraph?

Formatted: Strikethrough

Formatted: Strikethrough

Commented [JL3]: I would actually remove all of this. The paper is too top heavy—lots of intro but not enough substance in the body of the paper

Formatted: Strikethrough

Formatted: Strikethrough

Commented [JL4]: Ok, this is when the body of the paper should start. Substantive info—not overviews

Application of COWS in Skilled Nursing and Rehabilitation Facilities

The implementation of the Clinical Opiate Withdrawal Scale (COWS) in skilled nursing and rehabilitation facilities has garnered significant attention, as these settings often serve as a critical juncture in the treatment journey for individuals grappling with opioid use disorder. Researchers have undertaken a multifaceted approach to comprehensively assess the scale's efficacy and impact, exploring various dimensions of its application within these specialized care environments.

One crucial aspect of this ~~endeavor~~ assessment has been the examination of patient demographics and sample selection criteria. ~~Studies such as Snow & Wynn (2018) have focused~~

Editing Example

on a diverse range of individuals, encompassing those aged 18 to 80 years, ensuring a comprehensive representation of the populations served by these facilities. This inclusive approach has facilitated a deeper understanding of the scale's versatility and potential to address the unique needs of patients from varying backgrounds and life stages (Shah & Huecker, 2022).

Commented [JL5]: Which facilities?

Commented [JL6]: And what did they find? Be specific

Commented [JL7]: This is an unclear paragraph. What is the main idea of this paragraph? Do you think you could state it more directly? It gets lost--

Editing Example

Qualitative interviews with healthcare providers have been a valuable source of insight, offering first-hand perspectives on the practical implementation of COWS within skilled nursing and rehabilitation settings. These interviews have shed light on the challenges and successes experienced by professionals on the frontlines of opioid withdrawal management, capturing their observations and perceptions regarding the scale's impact on patient care (Dunn et al., 2020).

Complementing these qualitative accounts are quantitative survey methodologies, which have yielded empirical data on the efficacy of COWS in recognizing withdrawal symptoms. These surveys have encompassed patient-reported experiences and clinician assessments, enabling researchers to analyze potential discrepancies and understand the withdrawal phenomenon holistically (Canamo & Tronco, 2019).

Case study reviews have further enriched the research landscape, examining specific clinical scenarios and treatment regimens in-depth. These studies have explored the application of COWS in complex cases, such as those involving co-occurring disorders like posttraumatic stress disorder or during transitional phases from one pharmacotherapy to another, like the transition from buprenorphine to extended-release naltrexone (Comer et al., 2020).

Moreover, comparative analyses have been conducted to evaluate the performance of facilities that utilize COWS against those that do not employ this standardized scale. These comparisons have offered valuable insights into COWS implementation's potential advantages and limitations, shedding light on its impact on patient outcomes and quality of care (Snow & Wynn, 2018).

Effectiveness of COWS in Improving Patient Outcomes

At the core of evaluating the Clinical Opiate Withdrawal Scale (COWS) lies the pivotal

Commented [JL8]: Are you going to discuss the findings?

Commented [LR9]: And...what salient conclusions have been made from these studies? I think you want to highlight what was learned from this qualitative research within these settings rather than the fact that these studies have occurred.

Commented [LR10]: Don't be afraid to cite certain studies and get more detailed about the results.

Commented [JL11]: I think what you are doing is giving an overview of the types of studies conducted but you need the detail of what these studies actually found in specific terms

Editing Example

question of its effectiveness in enhancing patient outcomes within skilled nursing and rehabilitation facilities. Through a synthesis of research findings, this chapter elucidates the scale's impact on critical aspects of care, including recognizing withdrawal symptoms, the timeliness and accuracy of interventions, and its influence on quality of life metrics and overall patient and provider satisfaction.

A recurring theme across multiple studies is the COWS scale's proficiency in facilitating the comprehensive identification of withdrawal symptoms. Researchers have consistently observed that the systematic approach of COWS enables healthcare providers to pinpoint a constellation of physiological and psychological indicators, encompassing restlessness, anxiety, sweating, and pupil dilation, among others (Dunn et al., 2020; Snow & Wynn, 2018). By capturing this broad spectrum of symptoms, the scale empowers clinicians to delineate the etiology of withdrawal, discern its temporal progression, and evaluate its correlation with patient outcomes, thereby informing tailored treatment strategies (Comer et al., 2020).

Furthermore, the utilization of COWS has been associated with enhanced timeliness and accuracy in delivering interventions. The scale's quantitative scoring system and objective nature facilitate a standardized approach to assessing withdrawal severity, ensuring uniformity in treatment decisions across various healthcare settings (Canamo & Tronco, 2019). This consistency, coupled with the scale's sensitivity to symptom fluctuations, enables healthcare providers to administer appropriate interventions promptly, mitigating the exacerbation of withdrawal symptoms and optimizing patient comfort and safety.

The impact of COWS on patient-reported quality-of-life metrics has been the subject of extensive inquiry. Studies have highlighted the scale's potential to bridge the gap between subjective patient experiences and observer-rated assessments, leading to more personalized and

Commented [JL12]: What do you mean by this?

Commented [JL13]: What exactly did they study—what specific results or conclusions did they draw

Commented [JL14]: Give more details about the studies

Editing Example

effective treatment strategies (Dunn et al., 2020). By incorporating patient self-reports alongside clinical observations, healthcare professionals can tailor interventions to individual needs, potentially enhancing satisfaction and overall well-being during the arduous journey of opioid withdrawal management.

Editing Example

Closely intertwined with quality of life considerations is the aspect of patient and provider satisfaction with the treatment process. Researchers have posited that the implementation of COWS may contribute to increased satisfaction by fostering a more aligned understanding between patients and providers, facilitating smoother transitions during critical phases like the shift from buprenorphine to extended-release naltrexone, and ultimately promoting a more responsive and conscientious therapeutic framework (Comer et al., 2020; Snow & Wynn, 2018).

As the body of evidence continues to evolve, recommendations for best practices in COWS implementation have emerged. These include comprehensive staff training, integration into electronic medical record systems, and active patient involvement in the assessment process, all of which are pivotal to optimizing the scale's utility and ensuring accurate assessments, effective management, and successful transitions to long-term treatment options (Canamo & Tronco, 2019).

Concurrently, future research directions have been identified, encompassing the need for methodologically rigorous randomized clinical trials, comparative assessments of patient-reported and observer-rated symptoms, and explorations of the relationship between withdrawal symptoms, as measured by COWS, and clinical outcomes such as treatment success, quality of life, and patient mortality (Dunn et al., 2020; Snow & Wynn, 2018). These endeavors hold the potential to solidify further the evidence base surrounding the Clinical Opiate Withdrawal Scale, refining its application and contributing to the ongoing evolution of addiction treatment approaches.

Review of Literature on COWS

The Clinical Opiate Withdrawal Scale (COWS) has emerged as a pivotal tool in the assessment and management of opioid withdrawal symptoms, particularly within skilled nursing

Commented [JL15]: Isn't this all a literature review?

Formatted: Centered, Indent: Left: 0"

Editing Example

and rehabilitation facilities. This section delves into the historical background, implementation mechanisms, and empirical evidence supporting the efficacy of COWS in enhancing patient care and outcomes.

- A. **Historical Background of COWS Usage:** The development of COWS can be traced back to the growing recognition ~~of for~~ the need for standardized and objective measures to evaluate the withdrawal process from opioids. Traditional methods relying solely on subjective patient reports or clinician observations were often prone to inconsistencies and biases (NIH, n.d). COWS emerged as a structured approach, combining observable physical symptoms with objective scoring criteria, enabling healthcare professionals to gauge the severity of withdrawal accurately and tailor treatment interventions accordingly.
- B. **Mechanism and Implementation of COWS:** COWS is an 11-item scale that encompasses a range of physical and psychological indicators associated with opioid withdrawal. These include factors such as pupillary dilation, gastrointestinal distress, muscle aches, and anxiety (NIH, n.d). By systematically evaluating and scoring each item, healthcare providers can quantify the overall withdrawal severity, facilitating data-driven decision-making regarding medication adjustments, supportive therapies, or environmental modifications. Implementing COWS within skilled nursing and rehabilitation facilities often involves staff training, protocol development, and integrating electronic health records (EHRs) to streamline documentation and information sharing (Jones et al., 2021).
- C. **Empirical Evidence Supporting COWS Effectiveness:** Numerous studies have underscored the clinical utility and efficacy of COWS in managing opioid withdrawal symptoms. Studies by Dunn et al. (2020) and Canamo and Tronco (2019) have delved deeper into the intricacies of opiate withdrawal assessment, while contributions from Snow and

Commented [JL16]: This info would fit better as an introduction to COWS in the Introduction section

Editing Example

Wynn (2018) and Comer et al. (2020) have expanded our knowledge of effective management strategies, particularly in the context of transitioning patients from buprenorphine to extended-release naltrexone. Research has demonstrated COWS' ability to enhance patient comfort, reduce the risk of complications, and improve treatment adherence (*Evaluating the Quality of Health Care*, n.d.). By offering a standardized and objective approach, COWS mitigates the subjectivity inherent in self-reported withdrawal assessments, ensuring consistent

Commented [JL17]: These studies should be discussed, fleshed out in further detail

Editing Example

and reliable monitoring of patient progress (Werder, 2015). Furthermore, integrating COWS into EHRs and remote patient monitoring solutions has facilitated real-time data acquisition, enabling prompt interventions and tailored care plans (Desai & Nadeski, 2016).

The literature review on COWS highlights its pivotal role in optimizing withdrawal management within skilled nursing and rehabilitation facilities. By providing a structured framework for symptom assessment and leveraging technological advancements, COWS has revolutionized the approach to opioid withdrawal treatment, laying the foundation for enhanced patient outcomes and quality of life.

Impact of COWS on Patient Quality of Life

The implementation of the Clinical Opiate Withdrawal Scale (COWS) within skilled nursing and rehabilitation facilities has far-reaching implications for patients' quality of life, extending beyond the immediate management of withdrawal symptoms. Its impact can be observed across various domains, including mental health, physical stability, and enhancing social relationships, all of which contribute to overall well-being and successful recovery.

A. **Mental** Health Outcomes: Opioid withdrawal can be an emotionally turbulent experience, often accompanied by anxiety, depression, and psychological distress (*Evaluating the Quality of Health Care*, n.d.). By providing a structured approach to symptom assessment and management, COWS equips healthcare providers with the tools necessary to mitigate the mental health challenges associated with withdrawal. By effectively alleviating physical discomfort, patients may experience reduced anxiety and more excellent stability, enabling them to focus more fully on their rehabilitation journey (Jones et al., 2021).

Furthermore, integrating COWS into patient-centered care models fosters increased engagement and shared decision-making. When patients actively understand their COWS scores and collaborate with providers on treatment plans, they can experience a heightened

Commented [JL18]: You keep reverting back to an introductory structure.

Commented [JL19]: I don't think these need to be lettered.

Editing Example

sense of autonomy and empowerment (Werder, 2015). This involvement can serve as a buffer against feelings of helplessness and depression, ultimately enhancing emotional well-being during the critical recovery phase.

B. Physical Stability Improvements: Effectively managing withdrawal symptoms through COWS improves patients' physical stability and comfort levels. By accurately assessing and addressing symptoms such as nausea, muscle aches, and gastrointestinal distress, healthcare providers can tailor medication regimens and supportive interventions, minimizing discomfort and promoting a more conducive environment for healing (NIH, n.d).

Moreover, integrating COWS into electronic health records (EHRs) and remote patient monitoring solutions enables real-time patient condition tracking, facilitating prompt interventions and adjustments to treatment plans as needed (Desai & Nadeski, 2016). This proactive approach enhances physical stability and fosters a sense of security and trust in the care provided, further contributing to overall well-being.


C. Enhancement of Social Relationships: Opioid withdrawal can strain interpersonal dynamics, potentially hindering an individual's ability to engage in meaningful social interactions and strengthen relationships (*Digital Transformation and the Patient Experience*, 2021). By effectively managing withdrawal symptoms through COWS, patients may experience improved physical and emotional capacity to participate in social activities and maintain connections with family and friends.

Furthermore, the objective assessment provided by COWS can facilitate the involvement of support systems and loved ones in the treatment process. By sharing withdrawal progress and collaborating on care plans, patients and their families can foster a deeper understanding and stronger bonds, ultimately contributing to successful reintegration into social circles and the

Commented [JL20]: Citation?

Editing Example

broader community.



Editing Example

The impact of COWS on patient quality of life within skilled nursing and rehabilitation facilities is multifaceted, transcending the immediate management of withdrawal symptoms. COWS catalyzes holistic well-being through its influence on mental health, physical stability, and social relationships, empowering patients to navigate the challenges of recovery with greater resilience and support.

Comparative Analysis and Future Directions

Commented [JL21]: This is the best section of the paper

While the Clinical Opiate Withdrawal Scale (COWS) has established itself as a valuable tool in the assessment and management of opioid withdrawal, it is essential to contextualize its effectiveness by contrasting it with alternative assessment methods. Additionally, acknowledging the challenges and limitations inherent in current research can pave the way for future studies that further refine and enhance the application of COWS in skilled nursing and rehabilitation facilities.

A. Comparison with Other Opioid Withdrawal Scales: The Subjective Opiate Withdrawal Scale (SOWS) and the Short Opiate Withdrawal Scale (SOWS-Gossop) represent alternative approaches to assessing opioid withdrawal symptoms (NIH, n.d). Unlike COWS, which primarily focuses on observable physical indicators, these scales heavily rely on patient self-reports of subjective experiences, such as cravings, anxiety, and perceived discomfort. While this subjective approach provides valuable insights into the patient's internal state, it may be influenced by individual pain thresholds, psychological factors, and potential response biases (*Evaluating the Quality of Health Care*, n.d.).

In contrast, COWS's emphasis on objective, measurable symptoms offers a more consistent and standardized assessment across diverse patient populations and healthcare settings. This objectivity can be particularly advantageous in skilled nursing and rehabilitation facilities, where consistency and reliability in symptom evaluation are paramount for effective treatment

Editing Example

planning and medication management (Jones et al., 2021).

However, it is crucial to recognize that a comprehensive approach to withdrawal assessment often benefits from including objective and subjective measures. While COWS excels in capturing the physical manifestations of withdrawal, the subjective scales can provide complementary insights into the patient's emotional and psychological experiences, which is equally essential in promoting holistic well-being (Werder, 2015).

B. Challenges and Limitations in Current Research: Despite the substantial body of evidence supporting the effectiveness of COWS, several challenges and limitations persist in current research. One notable concern is the potential for inter-rater variability, where different healthcare professionals may interpret and score withdrawal symptoms differently, leading to inconsistencies in assessment (*Evaluating the Quality of Health Care*, n.d.). Rigorous staff training and ongoing quality assurance measures are essential to mitigate this challenge.

Furthermore, while COWS provides a systematic framework for assessing withdrawal severity, it is crucial to consider individual patient factors, such as pre-existing health conditions, comorbidities, and personal experiences, which may influence the manifestation and perception of withdrawal symptoms. ~~Please account for these~~ These individual variables should be accounted for to avoid suboptimal treatment plans or inaccurate interpretations of patient outcomes (Jones et al., 2021).

C. Recommendations for Future Studies: Several avenues for future research can be explored to address the limitations and further enhance the application of COWS in skilled nursing and rehabilitation facilities. Longitudinal studies examining the long-term impact of COWS-guided withdrawal management on patient quality-of-life indicators would be invaluable in assessing the scale's broader efficacy beyond the immediate withdrawal phase.

Editing Example

Additionally, research into integrating COWS with emerging technologies, such as wearable devices and remote patient monitoring systems, could unlock new possibilities for continuous, real-time assessment and intervention, potentially improving patient outcomes and reducing the burden on healthcare staff (Desai & Nadeski, 2016).

Editing Example

Finally, exploring the potential for machine learning and artificial intelligence to enhance the interpretation and application of COWS data could lead to more personalized and adaptive treatment plans, further optimizing patient care in skilled nursing and rehabilitation facilities (*Digital Transformation and the Patient Experience*, 2021).

As the healthcare field continues to evolve, embracing innovative approaches and leveraging emerging technologies will be crucial in refining the use of COWS and ensuring its continued relevance in the pursuit of improved patient quality of life and overall well-being.

Conclusion

The Clinical Opiate Withdrawal Scale (COWS) has emerged as a transformative tool in the management of opioid withdrawal, particularly within skilled nursing and rehabilitation facilities. This traditional narrative literature review has comprehensively explored COWS's effectiveness in enhancing patient quality of life, delving into its historical background, implementation mechanisms, and empirical evidence supporting its efficacy.

The findings presented in this review underscore the multifaceted impact of COWS on various aspects of patient well-being, including mental health, physical stability, and social relationships. By providing a structured and objective approach to withdrawal symptom assessment, COWS empowers healthcare providers to deliver tailored interventions that alleviate discomfort, mitigate psychological distress, and foster a supportive environment conducive to recovery (NIH, n.d) (*Evaluating the Quality of Health Care*, n.d.).

The integration of COWS into electronic health records (EHRs) and the adoption of health information technology (HIT) have further amplified its potential, enabling real-time monitoring, prompt interventions, and enhanced patient engagement (Werder, 2015). This synergy between clinical expertise and technological advancement exemplifies the commitment of skilled nursing and rehabilitation facilities to deliver patient-centered care that prioritizes

Editing Example

quality of life-outcomes.

While the existing literature consistently highlights the benefits of COWS in withdrawal management, this review has also addressed the challenges and limitations that warrant further investigation. Notably, the potential for inter-rater variability and the need to consider individual patient factors underscore the importance of rigorous staff training and continuous quality improvement initiatives (Jones et al., 2021).

As the healthcare landscape continues to evolve, future research should explore the integration of COWS with emerging technologies, such as wearable devices, remote patient monitoring systems, and artificial intelligence. These advancements unlock new avenues for personalized care, adaptive treatment plans, and real-time interventions, further optimizing patient outcomes and quality of life (Desai & Nadeski, 2016); (*Digital Transformation and the Patient Experience*, 2021).

In conclusion, the Clinical Opiate Withdrawal Scale (COWS) represents a pivotal advancement in pursuing holistic patient care within skilled nursing and rehabilitation facilities. By facilitating the effective management of opioid withdrawal symptoms, COWS not only enhances immediate symptom relief but also paves the way for long-term improvements in mental health, physical stability, and social engagement. As healthcare providers and researchers continue to refine and expand the application of COWS, the ultimate beneficiaries will be the individuals navigating the complex recovery journey, empowered to reclaim their well-being and embrace a higher quality of life.

References

Desai, P., & Nadeski, M. (2016). *Remote Patient Monitoring Solutions*.
<https://www.ti.com/lit/pdf/ssztba9>

Digital Transformation and the Patient Experience. (2021). Healthcare Communications.

<https://healthcare-communications.com/wp-content/uploads/2021/12/Digital-Transformation-the-Patient-Experience-Whitepaper.pdf>

Commented [JL22]: Needs to be completed

Formatted: English (United States)

Formatted: English (United States)

Editing Example

Dunn, K., Bergeria, C., Huhn, A. S., & Strain, E. C. (2020). Differences in patient-reported and observer-rated opioid withdrawal symptom etiology, time course, and relationship to clinical outcome. *Drug and alcohol dependence*, 215, 108212.

<https://doi.org/10.1016/j.drugalcdep.2020.108212> *Evaluating the Quality of Health Care* (n.d.).

<https://obssr.od.nih.gov/sites/obssr/files/Evaluating-the-Quality-of-Health-Care.pdf>

Jones, B., Kwong, E., & Warburton, W. (2021). Quality Improvement Made Simple What Everyone Should Know about Health Care Quality Improvement Quick Guide. *Quality Improvement Made Simple What Everyone Should Know about Health Care Quality Improvement*, 3(1).

<https://www.health.org.uk/sites/default/files/QualityImprovementMadeSimple.pdf>

Kosten, T. R., & Baxter, L. E. (2019). Review article: Effective management of opioid withdrawal symptoms: A gateway to opioid dependence treatment. *The American journal on addictions*, 28(2), 55–62. <https://doi.org/10.1111/ajad.12862>

NIH. (n.d.). Clinical opiate withdrawal scale. Clinical Opioid Withdrawal Scale.

<https://nida.nih.gov/sites/default/files/ClinicalOpiateWithdrawalScale.pdf>

Nuamah, J. K., Sasangohar, F., Erraguntla, M., & Mehta, R. K. (2019). The past, present and future of opioid withdrawal assessment: a scoping review of scales and technologies. *BMC medical informatics and decision making*, 19(1), 113. <https://doi.org/10.1186/s12911-019-0834-8>

Werder, M. (2015). Health Information Technology: A key ingredient of the patient experience. *Patient Experience Journal*, 2(1), 143–147. <https://doi.org/10.35680/2372-0247.1071>

Formatted: English (United States)