Subject: AI, prepare to talk about your asthma.
Date: Wednesday, October 22, 2025 at 10:46:03 AM Eastern Daylight Time
From: GSK
To: sciwriter@mac.com

View on web



TRELEGY ELLIPTA
(fluticasone furoate 100 mcg, umeclidinium 62.5 mcg, and vilanterol 25 mcg inhalation powder)

Prescribing Information
Patient Information

About TRELEGY

TRELEGY is a prescription medicine used long term to treat asthma in adults. TRELEGY is not used to relieve sudden breathing problems and won't replace a rescue inhaler.

Important Safety Information

TRELEGY contains vilanterol. Long-acting beta$_2$-adrenergic agonist (LABA) medicines such as vilanterol, when used alone, increase the risk of hospitalizations and death from asthma problems. TRELEGY contains an inhaled corticosteroid (ICS), an anticholinergic, and a LABA. When an ICS and LABA are used together, there is **not** a significant increased risk in hospitalizations and death from asthma problems. **Continued below** ▼




When it comes to your asthma, be prepared.

It helps to organize your thinking if you write everything down for your appointment.

You may want to include information such as:

- **A record of symptoms you experience regularly**
- **Notes on what you believe could be triggering your symptoms**
- **A list of medicines you're currently taking**
- **Copies of medical documents from other doctors, emergency rooms, or urgent care—if available**
- **A list of relevant questions**

No matter if your visit is in person, virtual, or by phone, being prepared and taking the time to have an informed conversation will give your doctor the information they need to help make decisions about your treatment.



An interactive guide for your next doctor visit.

Not sure how to start a conversation with your doctor? We're here to help. Use our interactive Doctor Discussion Guide to help get the most from your next doctor's appointment.

GET THE GUIDE

Have you already been prescribed TRELEGY?

YES

NO

Let us know so we can make sure to provide you with the right support, such as savings coupons, tips on starting a routine, and more!

Important Safety Information (cont'd)

- **Do not use TRELEGY to relieve sudden breathing problems.** Always have a rescue inhaler with you to treat sudden symptoms.
- **TRELEGY should not be used in children younger than 18 years of age.** It is not known if TRELEGY is safe and effective in children.
- **Do not use TRELEGY** if you have a severe allergy to milk proteins or are allergic to any of the ingredients in TRELEGY. Ask your healthcare provider if you are not sure.
- **Do not** use TRELEGY more often than prescribed.
- **Do not take TRELEGY with other medicines that contain a LABA or an anticholinergic for any reason. Tell your healthcare provider about all your medical conditions and about all the medicines you take.**
- Call your healthcare provider or get medical care right away if your breathing problems get worse, if you need your rescue inhaler more often than usual or it does not work as well to relieve your symptoms.
- **TRELEGY can cause serious side effects, including:**
  - **fungal infection in your mouth or throat (thrush).** Rinse your mouth with water without swallowing after using TRELEGY to help reduce your chance of getting thrush.
  - **weakened immune system and increased chance of getting infections.**
  - **reduced adrenal function.** This can happen when you stop taking oral corticosteroids and start taking inhaled corticosteroids.
  - **sudden breathing problems immediately after inhaling TRELEGY.** If this happens, stop taking TRELEGY and call your healthcare provider right away.
  - **serious allergic reactions.** Call your healthcare provider or get emergency medical care if you get any of the following symptoms: rash; hives; swelling of your face, mouth, and tongue; breathing problems.
  - **effects on heart:** increased blood pressure; a fast or irregular heartbeat, awareness of heartbeat; chest pain.
  - **effects on nervous system:** tremor; nervousness.
  - **bone thinning or weakness (osteoporosis).**
  - **eye problems** including glaucoma, increased pressure in your eye, cataracts, blurred vision, worsening of narrow-angle glaucoma, or other changes in vision. You should have regular eye exams while using TRELEGY. Acute narrow-angle glaucoma can cause permanent loss of vision if not treated. Symptoms of acute narrow-angle

glaucoma may include: eye pain or discomfort; nausea or vomiting; blurred vision; seeing halos or bright colors around lights; red eyes. If you have these symptoms, call your healthcare provider right away before taking another dose.

- **new or worse urinary retention.** Symptoms may include: difficult, painful, or frequent urination; urination in a weak stream or drips. If you have these symptoms, stop taking TRELEGY and call your healthcare provider right away.
- **changes in laboratory blood values,** including high levels of blood sugar (hyperglycemia) and low levels of potassium (hypokalemia).
- **slowed growth in children.**

- **Common side effects of TRELEGY include:** runny nose and sore throat; upper respiratory tract infection; bronchitis; respiratory tract infection; inflammation of the sinuses; painful and frequent urination (signs of a urinary tract infection); flu; headache; back pain.

Please see full Prescribing Information, including Patient Information, for TRELEGY.

GSK Help Center

Explore the GSK Help Center website to find customer service support and contact information.
Contact Us

GSK for You

If you have questions about how to access your GSK medication or need cost information, visit gskforyou.com or call 1-866-GSK-FOR-U (1-866-475-3678).

This email is intended for US residents only.

STOP OR CHANGE EMAIL COMMUNICATION: If you wish to stop receiving these email communications, or wish to stop receiving all email from GSK, Unsubscribe.

**You are encouraged to report negative side effects of prescription drugs to the FDA.
Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.**

The shape of the ELLIPTA inhaler is a trademark of the GSK group of companies.

Trademarks are owned by or licensed to the GSK group of companies.

GSK

©2025 GSK or licensor. All rights reserved.
Produced in USA.

GSK, 410 Blackwell Street, Durham, NC 27701



Privacy Notice | Contact Us

Please do not respond to this message. Mail sent to this email address cannot be answered.

PMUS-FVUEML250091 July 2025