

Subject: Al, it's time to focus on your breathing.

Date: Thursday, October 9, 2025 at 10:47:30 AM Eastern Daylight Time

From: GSK

To: sciwriter@mac.com

[View on web](#)



TRELEGY ELLIPTA
(fluticasone furoate 100 mcg, umecidinium 62.5 mcg,
and vilanterol 25 mcg inhalation powder)

[Prescribing Information](#)
[Patient Information](#)

About TRELEGY

TRELEGY is a prescription medicine used long term to treat asthma in adults. TRELEGY is not used to relieve sudden breathing problems and won't replace a rescue inhaler.

Important Safety Information

TRELEGY contains vilanterol. Long-acting beta₂-adrenergic agonist (LABA) medicines such as vilanterol, when used alone, increase the risk of hospitalizations and death from asthma problems. TRELEGY contains an inhaled corticosteroid (ICS), an anticholinergic, and a LABA. When an ICS and LABA are used together, there is **not** a significant increased risk in hospitalizations and death from asthma problems. [Continued below ▼](#)

**NEED A DOCTOR
WHO UNDERSTANDS
YOUR ASTHMA?**

FIND A DOCTOR



When your asthma isn't under control, it can lead to difficulty breathing, impaired lung function, flare-ups, and frequent or recurring symptoms that affect your daily life.

Whether you think you may have asthma or you've been diagnosed, you can find a healthcare professional to help you.

UP TO 5X GREATER

Get up to 5X greater lung function improvement with TRELEGY.*

With 3 medicines in 1 inhaler, TRELEGY improves lung function so you can breathe more freely all day and night.

Your results may vary.
TRELEGY does not replace a rescue inhaler.

*Compared to those taking 2 of 3 medicines in TRELEGY, patients taking low-dose TRELEGY had an improvement in FEV₁ of 134 mL vs 24 mL, while patients taking high-dose TRELEGY had an improvement in FEV₁ of 168 mL vs 76 mL. FEV stands for forced expiratory volume. FEV₁ is the amount of air a person can force from their lungs in 1 second.

**SEE IF TRELEGY IS
RIGHT FOR YOU**

What should you say to your doctor?

You can use this handy discussion guide to help you get the most from a visit to your doctor.

GET THE GUIDE

What's next? Schedule an office or telemedicine visit. Ask your doctor about trying once-daily TRELEGY.

Important Safety Information (cont'd)

- **Do not use TRELEGY to relieve sudden breathing problems.** Always have a rescue inhaler with you to treat sudden symptoms.
- **TRELEGY should not be used in children younger than 18 years of age.** It is not known if TRELEGY is safe and effective in children.
- **Do not use TRELEGY** if you have a severe allergy to milk proteins or are allergic to any of the ingredients in TRELEGY. Ask your healthcare provider if you are not sure.

- **Do not** use TRELEGY more often than prescribed.
- **Do not take TRELEGY with other medicines that contain a LABA or an anticholinergic for any reason. Tell your healthcare provider about all your medical conditions and about all the medicines you take.**
- Call your healthcare provider or get medical care right away if your breathing problems get worse, if you need your rescue inhaler more often than usual or it does not work as well to relieve your symptoms.
- **TRELEGY can cause serious side effects, including:**
 - **fungal infection in your mouth or throat (thrush).** Rinse your mouth with water without swallowing after using TRELEGY to help reduce your chance of getting thrush.
 - **weakened immune system and increased chance of getting infections.**
 - **reduced adrenal function.** This can happen when you stop taking oral corticosteroids and start taking inhaled corticosteroids.
 - **sudden breathing problems immediately after inhaling TRELEGY.** If this happens, stop taking TRELEGY and call your healthcare provider right away.
 - **serious allergic reactions.** Call your healthcare provider or get emergency medical care if you get any of the following symptoms: rash; hives; swelling of your face, mouth, and tongue; breathing problems.
 - **effects on heart:** increased blood pressure; a fast or irregular heartbeat, awareness of heartbeat; chest pain.
 - **effects on nervous system:** tremor; nervousness.
 - **bone thinning or weakness (osteoporosis).**
 - **eye problems** including glaucoma, increased pressure in your eye, cataracts, blurred vision, worsening of narrow-angle glaucoma, or other changes in vision. You should have regular eye exams while using TRELEGY. Acute narrow-angle glaucoma can cause permanent loss of vision if not treated. Symptoms of acute narrow-angle glaucoma may include: eye pain or discomfort; nausea or vomiting; blurred vision; seeing halos or bright colors around lights; red eyes. If you have these symptoms, call your healthcare provider right away before taking another dose.
 - **new or worse urinary retention.** Symptoms may include: difficult, painful, or frequent urination; urination in a weak stream or drips. If you have these symptoms, stop taking TRELEGY and call your healthcare provider right away.
 - **changes in laboratory blood values,** including high levels of blood sugar (hyperglycemia) and low levels of potassium (hypokalemia).
 - **slowed growth in children.**
- **Common side effects of TRELEGY include:** runny nose and sore throat; upper respiratory tract infection; bronchitis; respiratory tract infection; inflammation of the sinuses; painful and frequent urination (signs of a urinary tract infection); flu; headache; back pain.

Please see full [Prescribing Information](#), including [Patient Information](#), for TRELEGY.

GSK Is Here to Help!
Contact Us



GSK *for you*

If you don't have prescription coverage,
or need help paying for your medicine,
visit [GSKforyou.com](https://www.gskforyou.com) or call
1-866-GSK-FOR-U (1-866-475-3678)

This email is intended for US residents only.

STOP OR CHANGE EMAIL COMMUNICATION: If you wish to stop receiving these email communications, or wish to stop receiving all email from GSK, [Unsubscribe](#).

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

The shape of the ELLIPTA inhaler is a trademark of the GSK group of companies.

Trademarks are owned by or licensed to the GSK group of companies.



©2024 GSK or licensor. All rights reserved. Produced in USA.

GSK, 410 Blackwell Street, Durham, NC 27701



[Privacy Notice](#) | [Legal Notices](#) | [Contact Us](#)

Please do not respond to this message. Mail sent to this email address cannot be answered.

PMUS-FVUEML240012 April 2024